

Table 22. Reported cases of HIV infection (not AIDS) for female adults and adolescents, by transmission category and race/ethnicity, 2006 and cumulative—45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting

| Transmission category | 2006 | | Cumulative ^a | |
|---|--------------|------------|-------------------------|------------|
| | No. | % | No. | % |
| White, not Hispanic | | | | |
| Injection drug use | 690 | 26 | 4,260 | 26 |
| Hemophilia/coagulation disorder | 3 | 0 | 24 | 0 |
| High-risk heterosexual contact ^b | 1,132 | 42 | 7,593 | 46 |
| Sex with injection drug user | 205 | 8 | 1,876 | 11 |
| Sex with bisexual male | 98 | 4 | 659 | 4 |
| Sex with person with hemophilia | 10 | 0 | 115 | 1 |
| Sex with HIV-infected transfusion recipient | 7 | 0 | 64 | 0 |
| Sex with HIV-infected person, risk factor not specified | 812 | 30 | 4,879 | 29 |
| Receipt of blood transfusion, blood components, or tissue | 12 | 0 | 146 | 1 |
| Other/risk factor not reported or identified | 850 | 32 | 4,567 | 28 |
| Total | 2,687 | 100 | 16,590 | 100 |
| Black, not Hispanic | | | | |
| Injection drug use | 1,034 | 13 | 7,105 | 14 |
| Hemophilia/coagulation disorder | 1 | 0 | 35 | 0 |
| High-risk heterosexual contact ^b | 3,451 | 44 | 23,732 | 46 |
| Sex with injection drug user | 313 | 4 | 3,473 | 7 |
| Sex with bisexual male | 155 | 2 | 1,294 | 3 |
| Sex with person with hemophilia | 5 | 0 | 71 | 0 |
| Sex with HIV-infected transfusion recipient | 14 | 0 | 120 | 0 |
| Sex with HIV-infected person, risk factor not specified | 2,964 | 38 | 18,774 | 36 |
| Receipt of blood transfusion, blood components, or tissue | 26 | 0 | 324 | 1 |
| Other/risk factor not reported or identified | 3,370 | 43 | 20,244 | 39 |
| Total | 7,882 | 100 | 51,440 | 100 |
| Hispanic | | | | |
| Injection drug use | 339 | 15 | 2,280 | 18 |
| Hemophilia/coagulation disorder | 2 | 0 | 11 | 0 |
| High-risk heterosexual contact ^b | 1,198 | 52 | 6,264 | 50 |
| Sex with injection drug user | 218 | 9 | 1,324 | 11 |
| Sex with bisexual male | 62 | 3 | 266 | 2 |
| Sex with person with hemophilia | 1 | 0 | 11 | 0 |
| Sex with HIV-infected transfusion recipient | 7 | 0 | 39 | 0 |
| Sex with HIV-infected person, risk factor not specified | 910 | 39 | 4,624 | 37 |
| Receipt of blood transfusion, blood components, or tissue | 12 | 1 | 58 | 0 |
| Other/risk factor not reported or identified | 767 | 33 | 3,843 | 31 |
| Total | 2,318 | 100 | 12,456 | 100 |

Table 22. Reported cases of HIV infection (not AIDS) for female adults and adolescents, by transmission category and race/ethnicity, 2006 and cumulative—45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting (*cont*)

| Transmission category | 2006 | | Cumulative ^a | |
|---|---------------------------|------------|---------------------------|------------|
| | No. | % | No. | % |
| Asian/Pacific Islander | | | | |
| Injection drug use | 10 | 7 | 23 | 5 |
| Hemophilia/coagulation disorder | 0 | 0 | 1 | 0 |
| High-risk heterosexual contact ^b | 57 | 42 | 198 | 42 |
| Sex with injection drug user | 7 | 5 | 18 | 4 |
| Sex with bisexual male | 8 | 6 | 15 | 3 |
| Sex with person with hemophilia | 0 | 0 | 0 | 0 |
| Sex with HIV-infected transfusion recipient | 0 | 0 | 0 | 0 |
| Sex with HIV-infected person, risk factor not specified | 42 | 31 | 165 | 35 |
| Receipt of blood transfusion, blood components, or tissue | 0 | 0 | 5 | 1 |
| Other/risk factor not reported or identified | 70 | 51 | 239 | 51 |
| Total | 137 | 100 | 466 | 100 |
| American Indian/Alaska Native | | | | |
| Injection drug use | 29 | 34 | 112 | 28 |
| Hemophilia/coagulation disorder | 0 | 0 | 0 | 0 |
| High-risk heterosexual contact ^b | 38 | 44 | 190 | 48 |
| Sex with injection drug user | 8 | 9 | 54 | 14 |
| Sex with bisexual male | 4 | 5 | 17 | 4 |
| Sex with person with hemophilia | 0 | 0 | 2 | 1 |
| Sex with HIV-infected transfusion recipient | 0 | 0 | 0 | 0 |
| Sex with HIV-infected person, risk factor not specified | 26 | 30 | 117 | 29 |
| Receipt of blood transfusion, blood components, or tissue | 0 | 0 | 1 | 0 |
| Other/risk factor not reported or identified | 19 | 22 | 94 | 24 |
| Total | 86 | 100 | 397 | 100 |
| Total cases | | | | |
| Injection drug use | 2,124 | 16 | 13,925 | 17 |
| Hemophilia/coagulation disorder | 6 | 0 | 73 | 0 |
| High-risk heterosexual contact ^b | 5,939 | 45 | 38,315 | 47 |
| Sex with injection drug user | 757 | 6 | 6,791 | 8 |
| Sex with bisexual male | 328 | 2 | 2,265 | 3 |
| Sex with person with hemophilia | 16 | 0 | 201 | 0 |
| Sex with HIV-infected transfusion recipient | 28 | 0 | 223 | 0 |
| Sex with HIV-infected person, risk factor not specified | 4,810 | 36 | 28,835 | 35 |
| Receipt of blood transfusion, blood components, or tissue | 50 | 0 | 536 | 1 |
| Other/risk factor not reported or identified | 5,106 | 39 | 29,312 | 36 |
| Total | 13,225^c | 100 | 82,161^d | 100 |

Note. See Table 16 for list of 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December 2006.

^a From the beginning of the epidemic through December 2006.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes 115 females of unknown race or multiple races.

^d Includes 812 females of unknown race or multiple races.